

CELESTIAL-TNCLL: An Ongoing, Open-Label, Multiregional, Phase 3 Study of Sonrotoclax (BGB-11417) + Zanubrutinib vs Venetoclax + Obinutuzumab for Treatment-Naive CLL

Antonio Cuneo,¹ Piers E.M. Patten,^{2,3} Arnon P. Kater,⁴ Jennifer A. Woyach,⁵ Jianyong Li,⁶ Talha Munir,⁷ Tommi Salmi,⁸ Patrick Phuong,⁹ Tian Tian,⁹ Mazyar Shadman^{10,11}

¹Hematology Section, St. Anna University Hospital, Ferrara, Italy; ²Comprehensive Cancer Centre, King's College London, London, UK; ³King's College Hospital, London, UK; ⁴Lymphoma and Myeloma Research Amsterdam, Amsterdam, the Netherlands; ⁵Ohio State University Comprehensive Cancer Center, Columbus, OH, USA; ⁶The First Affiliated Hospital of Nanjing Medical University, Jiangsu Province Hospital, Nanjing, China; ⁷Leeds Teaching Hospitals NHS Trust, Leeds, UK; ⁸BeiGene International GmbH, Basel, Switzerland; ⁹BeiGene USA, Inc, San Mateo, CA, USA; ¹⁰Fred Hutchinson Cancer Research Center, Seattle, WA, USA; ¹¹University of Washington, Seattle, WA, USA


INTRODUCTION

- The combination of venetoclax, the first-generation BCL2 inhibitor, and ibrutinib, a BTK inhibitor, has demonstrated efficacy in patients with CLL¹
- However, the toxicity profile of this regimen suggests a need for a more tolerable BTK/BCL2 inhibitor combination
- Sonrotoclax, a next-generation BCL2 inhibitor, is a more selective and more pharmacologically potent inhibitor of BCL2 than venetoclax²
- Zanubrutinib, a next-generation BTK inhibitor, significantly improved PFS and had a more tolerable safety profile, including fewer cardiac AEs, vs ibrutinib in a randomized, head-to-head study of patients with CLL/SLL³
- In a phase 1 study in patients with treatment-naive (TN) CLL, treatment with sonrotoclax + zanubrutinib was generally well-tolerated, with low rates of TEAEs, and resulted in high ORRs and deep responses⁴
- Presented here is the study design of CELESTIAL-TNCLL, a phase 3 trial of sonrotoclax + zanubrutinib in TN CLL

STUDY STATUS

- Enrollment for CELESTIAL-TNCLL began in December 2023, and the study is currently recruiting
- Approximately 230 study sites in 20 countries are planned (**Figure 2**), with an estimated enrollment of 640 patients. In the Americas, there are approximately 50 sites in the US, 6 in Brazil and 15 in Canada


Figure 2. CELESTIAL-TNCLL Study Sites (Planned)



METHODS

- CELESTIAL-TNCLL (BGB-11417-301; NCT06073821) is a randomized, open-label, phase 3 study comparing the efficacy of sonrotoclax + zanubrutinib vs venetoclax + obinutuzumab in patients with TN CLL (**Figure 1**)
- Efficacy will be assessed in accordance with 2018 iwCLL guidelines⁵ with modification of treatment-related lymphocytosis for patients with CLL⁶

Figure 1. CELESTIAL-TNCLL Study Design



^a Defined as CR or CR with incomplete recovery. ^b At <10⁻⁴ sensitivity at the first post-treatment follow-up based on next-generation sequencing by clonoSEQ[®] and flow cytometry. BM, bone marrow; CRR, complete response rate; DOR, duration of response; INV, assessed by investigator; IRC, assessed by independent review committee; PB, peripheral blood; R, randomized; TN, treatment naive; uMRD4, undetectable measurable residual disease.

REFERENCES

1. Wierda WG, et al. *J Clin Oncol*. 2021;29:3853-3865.
2. Hu N, et al. ACR 2020. Abstract 3077.
3. Brown JR, et al. *N Engl J Med*. 2023;388:319-332.
4. Tam CS, et al. ASH 2023. Abstract 327.
5. Hallek M, et al. *Blood*. 2018;131:2745-2760.
6. Cheson BD, et al. *J Clin Oncol*. 2012;30:2820-2822.

DISCLOSURES

AC: Fees for participation at congresses and advisory boards: AbbVie, AstraZeneca, BeiGene, Janssen, and Lilly. **PEMP:** Research funding: AstraZeneca, AbbVie, BeiGene, Janssen, Novartis; Honoraria: AstraZeneca, AbbVie, BeiGene; Travel, accommodations, expenses: AbbVie, BeiGene, Janssen. **APK:** Research funding: Janssen, Roche, Genentech, AbbVie, AstraZeneca, BMS; Honoraria: AbbVie; Consulting role: Janssen, AbbVie, Genentech, AstraZeneca, BMS, Lava Therapeutics; Travel, accommodations, expenses: AbbVie, Janssen. **JAW:** Consulting: AbbVie, AstraZeneca, BeiGene, Genentech, LOXO@Lilly, Merck, Newave, Pharmacyclics; Research funding: AbbVie, Janssen, BeiGene, Pharmacyclics. **JL:** Nothing to disclose. **TM:** Honoraria: BeiGene, AstraZeneca, Sobi, Roche, Janssen, AbbVie, Lilly; Consulting: AbbVie, BeiGene, Sobi, Alexion, Novartis, Janssen, AstraZeneca, Lilly, Roche; Research grants: Janssen, AbbVie; Travel, accommodations, expenses: Alexion, BeiGene, AbbVie, Janssen, AstraZeneca; Advisory board: AbbVie, BeiGene, AstraZeneca, Janssen. **TS:** Employment: BeiGene Switzerland GmbH; Equity holder: BeiGene Ltd. **PP, TT:** Employment: BeiGene. **MS:** Consultancy: AbbVie, Genentech, AstraZeneca, Genmab, Janssen, BeiGene, BMS, MorphoSys/Incyte, Kite Pharma, Eli Lilly, Fate Therapeutics, Nurix, Merck; Research funding: Mustang Bio, Genentech, AbbVie, BeiGene, AstraZeneca, Genmab, MorphoSys/Incyte, Vincerox; Stock options: Koi Biotherapeutics; Employment: BMS (spouse).

ACKNOWLEDGMENTS

The authors thank the patients and their families, investigators, co-investigators, and the study teams at each of the participating centers. This study was sponsored by BeiGene, Ltd. Medical writing support was provided by Angela R. Eder, PhD, of Nucleus Global, an Inizio company, and supported by BeiGene.