

## **Preliminary safety and antileukemic activity of sonrotoclax (BGB-11417), a potent and selective BCL2 inhibitor, in treatment-naive patients with acute myeloid leukemia**

**Authors:** Jake Shortt,<sup>1</sup> Shuh Ying Tan,<sup>2</sup> Sophie Leitch,<sup>3</sup> James Mao-Chun Liang,<sup>4</sup> Chun Yew Fong,<sup>5</sup> Sundra Ramanathan,<sup>6</sup> Qiubai Li,<sup>7</sup> Pau Montesinos,<sup>8</sup> Courtney DiNardo,<sup>9</sup> Jinhua Zhong,<sup>10</sup> Wai Y. Chan,<sup>11</sup> Rui Li,<sup>10</sup> Oleg Zernovak,<sup>11</sup> Amit Agarwal,<sup>11</sup> Andrew H. Wei<sup>12</sup>

**Affiliations:** <sup>1</sup>Monash Health and Monash University, Clayton, VIC, Australia; <sup>2</sup>St Vincent's Hospital, Melbourne, VIC, Australia; <sup>3</sup>Te Whatu Ora, Health New Zealand, Waitemata, Auckland, New Zealand; <sup>4</sup>Middlemore Hospital, Auckland, New Zealand; <sup>5</sup>Austin Health, Melbourne, VIC, Australia; <sup>6</sup>St George Hospital, Kogarah, NSW, Australia; <sup>7</sup>Union Hospital, Tongji Medical College, Huazhong University of Science and Technology, Wuhan, China; <sup>8</sup>Hospital Universitari I Politècnic La Fe, Valencia, Spain; <sup>9</sup>MD Anderson Cancer Center, Houston, TX, USA; <sup>10</sup>BeiGene (Shanghai) Co, Ltd, Shanghai, China; <sup>11</sup>BeiGene USA, Inc, San Mateo, CA, USA; <sup>12</sup>Peter MacCallum Cancer Centre, Royal Melbourne Hospital, Melbourne, VIC, Australia

### **ABSTRACT**

**Objectives:** B-cell lymphoma 2 (BCL2), a key regulator of apoptosis, is overexpressed in many hematologic malignancies. The BCL2 inhibitor venetoclax + azacitidine (AZA) has improved outcomes for treatment-naive (TN) patients with newly diagnosed acute myeloid leukemia (AML) unfit for intensive chemotherapy (TN unfit AML), with median overall survival (OS) of 14.7 months and complete remission in 36.7%. However, primary or secondary resistance is common, and long-term survival is suboptimal. Sonrotoclax (BGB-11417), a next-generation BCL2 inhibitor, is a selective and pharmacologically potent inhibitor of BCL2. In ongoing phase 1 studies, sonrotoclax has been well tolerated with preliminary antitumor activity in B-cell malignancies, AML/myelodysplastic syndrome (MDS), and multiple myeloma. Presented here are preliminary safety and antileukemic activity of sonrotoclax + AZA in TN unfit AML patients in BGB-11417-103 (NCT04771130; EudraCT: 2021-003285-12).

**Methods:** BGB-11417-103 is an ongoing, phase 1b/2, global, multicenter, dose finding and expansion study evaluating sonrotoclax + AZA in patients with AML (TN unfit [ $\geq 65$  years old or with comorbidities] or relapsed/refractory [R/R]), MDS, or MDS/myeloproliferative neoplasm. Among TN unfit AML patients, dose cohorts were evaluated to explore sonrotoclax 40 mg x 10 days, 80 mg x 10 days, 160 mg x 10 days, 160 mg x 28 days, 320 mg x 21 days, or 320 mg x 14 days + AZA (75 mg/m<sup>2</sup> x 7 days). For cycle 1, a 4-day sonrotoclax ramp-up was incorporated starting at one-eighth the target dose. Dose-limiting toxicities (DLTs) were assessed up to day 28 for nonhematologic toxicities and day 42 for hematologic toxicities (unless cycle 2 was initiated). Treatment-emergent adverse events (TEAEs) were graded per CTCAE v5.0. Response assessment was conducted per European Leukemia Net 2017 criteria.

**Results:** As of September 25, 2023, 42 enrolled TN patients with AML received study drug across dose escalation and expansion cohorts and were included in efficacy analyses; 16 (38%) remain on treatment. Median age was 75 years and median follow-up was 9.6 months. All patients had  $\geq 1$  TEAE. The most common grade  $\geq 3$  nonhematologic TEAEs were pneumonia (17%) and hypokalemia (14%); common grade  $\geq 3$  hematologic TEAEs were neutropenia (64%), thrombocytopenia (48%), febrile neutropenia (48%), and anemia (43%). Grade  $\geq 3$  infections occurred in 52% of patients. The most common TEAE leading to dose reduction was neutropenia (n=4, sonrotoclax reduction; n=9, AZA reduction). The most common TEAE class leading to sonrotoclax discontinuation was infection (n=5). No deaths due to treatment-related TEAEs or clinical tumor lysis syndrome (TLS) occurred; 1 patient had laboratory TLS

(160 mg x 10 days, cycle 2, hyperuricemia and hyperphosphatemia, resolved in 4 days). Mortality rate within 30 days was 2% (1 patient). CR and CR/CRh rates were 50% and 62%, with median time to response of 1.7 and 1.3 months, respectively; median duration of response for both was 18.8 months. Preliminary median OS was 20.6 (95% CI, 9.5-NE) months, with 64% OS at 12 months.

**Conclusions:** In this phase 1b/2 dose escalation and expansion study, sonrotoclax + AZA was generally well tolerated with promising antileukemic activity in TN unfit AML patients. Efficacy was seen starting at the lowest dose cohort. The study is ongoing in escalation, and the recommended phase 2 dose is still being determined.