

Combination Treatment With Novel BCL2 Inhibitor Sonrotoclax (BGB-11417) and Zanubrutinib Induces High Rate of Complete Remission for Patients With Relapsed/Refractory Mantle Cell Lymphoma

Constantine S. Tam,¹ Stephen S. Opat,² Marc S. Hoffmann,³ Jacob D. Soumerai,⁴ Masa Lasica,⁵ Narendranath Epperla,⁶ Jing-Zhou Hou,⁷ Ramón García-Sanz,⁸ Johannes Schetelig,⁹ Robert Weinkove,^{10,11} Yiqian Fang,¹² Sheel Patel,¹³ Wei Ding,¹³ Haiyi Guo,¹² Raul Cordoba¹⁴

¹Alfred Hospital and Monash University, Melbourne, VIC, Australia; ²Lymphoma Research Group, School of Clinical Sciences at Monash Health, Monash University, Clayton, VIC, Australia; ³University of Kansas Cancer Center, Kansas City, KS, USA; ⁴Massachusetts General Hospital Cancer Center and Harvard Medical School, Boston, MA, USA; ⁵St Vincent's Hospital Melbourne, Fitzroy, VIC, Australia; ⁶The James Cancer Hospital and Solove Research Institute at Ohio State University, Columbus, OH, USA; ⁷University of Pittsburgh Medical Center, Pittsburgh, PA, USA; ⁸Hospital Universitario de Salamanca, Salamanca, Spain; ⁹Universitätsklinikum Carl Gustav Carus An Der Technischen Universität Dresden, Dresden, Germany; ¹⁰Te Rerenga Ora Blood and Cancer Centre, Te Whatu Ora Health New Zealand Capital Coast & Hutt Valley, Wellington, New Zealand; ¹¹Cancer Immunotherapy Programme, Malaghan Institute of Medical Research, Wellington, New Zealand; ¹²BeiGene (Shanghai) Co, Ltd, Shanghai, China; ¹³BeiGene USA, Inc, San Mateo, CA, USA; ¹⁴Hospital Universitario Fundación Jiménez Díaz, Madrid, Spain

Presenter: Zhiyu Liang



Disclosure for Zhiyu Liang

- Employment and may own stock: BeiGene, Inc.

Introduction

- Combining BCL2 and BTK inhibition with venetoclax + ibrutinib has shown efficacy in patients with R/R MCL; however, this treatment was associated with high rates of toxicity and a need for a safer and potent combination still remains¹
- Sonrotoclax (sonro; BGB-11417), a next-generation BCL2 inhibitor, is a more selective and pharmacologically potent inhibitor of BCL2 than venetoclax with a shorter half-life and no accumulation²
- Zanubrutinib (zanu) is a next-generation BTK inhibitor approved in multiple (5) indications, including R/R MCL³
- Zanu was designed to provide complete and sustained BTK occupancy for efficacy across multiple B-cell malignancies with fewer off-target AEs compared with other BTK inhibitors^{4,5}
- Here, safety and efficacy data are presented for patients with R/R MCL treated with sonro + zanu in the ongoing BGB-11417-101 study


MCL, mantle cell lymphoma; R/R, relapsed/refractory.

1. Wang M, et al. ASH 2023. Abstract LBA-2; 2. Hu N, et al. AACR 2020. Abstract 3077; 3. Brukinsa. Prescribing information. BeiGene, Ltd; 2024; 4. Guo Y, et al. *J Med Chem*. 2019;62(17):7923-7940; 5. Tam CS, et al. *Expert Rev Clin Pharmacol*. 2021;14(11):1329-1344.



Study Design


- BGB-11417-101 (NCT04277637): first-in-human, phase 1, open-label, multicenter, dose-escalation and -expansion study in patients with B-cell malignancies
- Eligible patients: R/R MCL (disease that relapsed after or was refractory to ≥ 1 prior systemic therapy) and required treatment in the opinion of the investigator
- Primary objectives: assess safety/tolerability, evaluate the ramp-up dosing schedule, define MTD, and determine the RP2D of sonro in combination with zanu



^a The safety monitoring committee reviewed dose-level cohort data before dose escalation. ^b Zanu was administered orally (320 mg QD or 160 mg BID) 8 to 12 weeks prior to sonro treatment; sonro was administered orally, QD, following a daily or weekly ramp-up schedule to mitigate potential risk of TLS

Patient Disposition

- As of February 4, 2024, a total of 40 patients with R/R MCL received sonro + zanu and 27 remained on study treatment
 - 10 patients (25%) discontinued sonro + zanu due to PD (n=5), patient withdrawal (n=1), and AEs (n=4; 1 was treatment-related [pneumonia])
 - 4 patients discontinued zanu due to PD during lead-in (n=3) and 1 patient discontinued from zanu only due to diarrhea
- The sonro 160- and 320-mg dose levels were chosen for expansion cohorts



Baseline Patient Characteristics

Characteristic	Sonro 80 mg + zanu (n=6)	Sonro 160 mg + zanu (n=13)	Sonro 320 mg + zanu (n=16)	Sonro 640 mg + zanu (n=5)	All (N=40)
Study follow-up, median (range), months	27.5 (3.9-29.6)	16.0 (1.0-25.7)	12.5 (0.4-18.8)	3.5 (2.2-8.6)	12.5 (0.4-29.6)
Age, median (range), years	60.0 (46-84)	69.0 (45-81)	69.0 (45-85)	71.0 (68-80)	68.5 (45-85)
Male sex, n (%)	5 (83)	11 (85)	7 (44)	3 (60)	26 (65)
ECOG performance status, n (%)					
0	3 (50)	8 (62)	4 (25)	3 (60)	18 (45)
1	2 (33)	5 (38)	12 (75)	2 (40)	21 (53)
Tumor bulk, n (%)					
LDi <10 and ≥5 cm	3 (50)	4 (31)	3 (19)	2 (40)	12 (30)
LDi ≥10 cm	1 (17)	2 (15)	3 (19)	0	6 (15)
Ki67 proliferation index, n (%)					
<30%	3 (50)	4 (31)	6 (38)	0	13 (33)
≥30%	2 (33)	2 (15)	4 (25)	2 (40)	10 (25)
Prior therapy					
No. of lines of prior systemic therapy, median (range)	1 (1-1)	1 (1-4)	1 (1-3)	1 (1-1)	1 (1-4)
No. of lines of prior systemic therapy, n (%)					
1	6 (100)	10 (77)	11 (69)	5 (100)	32 (80)
2	0	2 (15)	1 (6)	0	3 (8)
≥3	0	1 (8)	4 (25)	0	5 (13)
Prior BTK inhibitor, n (%)	0	0	3 (19)	0	3 (8)^a
BTK inhibitor as last prior therapy, n (%)	0	0	3 (19)	0	3 (8)^a
Prior BTK inhibitor duration, median (range), months	–	–	4.8 (0.3-25.0)	–	4.8 (0.3-25.0)
Prior cellular therapies (transplant or CAR-T), n (%)	2 (33)	3 (23)	6 (38)	0	11 (28)

^a Two patients discontinued due to toxicity.
CAR-T, chimeric antigen receptor T-cell; LDi, longest diameter.

TEAE Summary

- Toxicity was generally the same among all tested dose levels with no new safety signals identified; sonro 160-mg and 320-mg dose levels were chosen for expansion cohorts

Patients, n (%)	Sonro 80 mg + zanu (n=6)	Sonro 160 mg + zanu (n=13)	Sonro 320 mg + zanu (n=16)	Sonro 640 mg + zanu (n=5)	All (N=40)
Any TEAE	4 (67)	13 (100)	15 (94)	5 (100)	37 (93)
Grade ≥3	4 (67)	6 (46)	7 (44)	1 (20)	18 (45)
Serious TEAEs	3 (50)	4 (31)	2 (13)	0	9 (23)
Leading to death	1 (17)	1 (8)	1 (6)	0	3 (8) ^a
Leading to zanu discontinuation	1 (17)	3 (23)	2 (13)	0	6 (15) ^b
Leading to zanu dose reduction	1 (17)	1 (8)	0	0	2 (5) ^c
Treated with sonro, n (%)	6 (100)	11 (85)	13 (81)	5 (100)	35 (88)
Leading to sonro discontinuation	0	3 (23)	2 (13)	0	5 (13) ^d
Leading to sonro dose reduction	0	0	0	0	0
Leading to death	0	1 (8)	0	0	1 (3) ^e


^a Pleural effusion (due to PD), abdominal sepsis, and pneumonia. ^b Lymph node pain (due to PD), diarrhea, MDS, abdominal sepsis, pneumonia, and bruising. ^c COVID-19 (temporary).

^d Diarrhea, abdominal sepsis, MDS, pneumonia and lymph node pain secondary to PD. ^e Pneumonia.

MDS, myelodysplastic syndrome; sonro, sonrotoclax; zanu, zanubrutinib.

TEAEs in ≥ 5 Patients by Grade


- Most common any-grade TEAEs: contusion (30%), neutropenia (28%), and diarrhea (28%)
- Most common grade ≥ 3 TEAE: neutropenia (18%)
 - Neutropenia was manageable, with:
 - No dose reductions
 - Only 1 dose hold due to concurrent COVID-19 infection
 - 6 patients used G-CSF (median duration = 3.5 days)
- No laboratory or clinical TLS
- Dose escalation completed with no MTD reached



^a Neutropenia combines preferred terms *neutrophil count decreased* and *neutropenia*. ^b Thrombocytopenia combines preferred terms *platelet count decreased* and *thrombocytopenia*.

Treatment Response Rates^a


- Median study follow-up was 12.5 months
 - ORRs were 73% and 92% in the 160- and 320-mg cohorts, respectively
 - CR rates were 46% and 83%, respectively



^a Responses were assessed per Lugano 2014 criteria. ^b ORR was defined as PR or better. ^c For all dose levels. ^d For all patients as treated (N=40).
 1. Cheson BD, et al. *J Clin Oncol*. 2014;32(27):3059-3068.

Treatment Duration and Investigator-Assessed Responses^a

- Of 3 response-evaluable patients with prior BTK inhibitor treatment, 2 responded: 1 PR and 1 CR



* Patient had prior treatment with BTK inhibitor.

^a Gray bar indicates duration of zanu lead-in.

Conclusions

- Sonrotoclax in combination with zanubrutinib was generally well tolerated
 - The maximum tolerated dose was not reached up to the highest assessed dose of 640 mg
 - No atrial fibrillation or TLS (laboratory or clinical) events were observed
- Sonrotoclax + zanubrutinib combination therapy demonstrated deep responses in patients with R/R MCL, including an ORR of 92% and CR rate of 83% in the 320-mg cohort
- The 320-mg dose was selected as RP2D for development in future pivotal studies

Acknowledgments

The authors thank the patients and their families, investigators, co-investigators, and the study teams at each of the participating centers. This study was sponsored by BeiGene, Ltd.

Medical writing was provided by Shanen Perumal, PhD, of Nucleus Global, an Inizio company, and supported by BeiGene.

Corresponding author:

Zhiyu Liang, zhiyu.liang@beigene.com