

Zanubrutinib in Acalabrutinib-Intolerant Patients With B-Cell Malignancies


Loïc Ysebaert,¹ Mazyar Shadman,² Ian W. Flinn,³ Edwin C. Kingsley,⁴ Benjamin Freeman,⁵ Moshe Y. Levy,⁶ Charles M. Farber,⁷ James D'Olimpio,⁸ Jennifer L. Cultrera,⁹ Ben Zhang,¹⁰ Rocco J. Crescenzo,¹¹ Adam Idoine,¹¹ Xiaoping Zhang,¹¹ Kunthel By,¹¹ Jeff P. Sharman¹²

¹Department of Hematology, University Cancer Institute of Toulouse Oncopole, Toulouse, France; ²Fred Hutchinson Cancer Research Center, University of Washington, Clinical Research Division, Seattle, WA, USA; ³Tennessee Oncology/OneOncology, Department of Medical Oncology, Nashville, TN, USA; ⁴Comprehensive Cancer Centers of Nevada, Department of Medical Oncology, Las Vegas, NV, USA; ⁵Summit Medical Group, Hematology and Medical Oncology, Florham Park, NJ, USA; ⁶Texas Oncology-Baylor Charles A. Sammons Cancer Center, Hematology and Medical Oncology, Dallas, TX, USA; ⁷Atlantic Hematology Oncology, Morristown Medical Center, Hematology and Medical Oncology, Morristown, NJ, USA; ⁸Clinical Research Alliance, Department of Clinical Research, Westbury, NY, USA; ⁹Florida Cancer Specialists & Research Institute, Hematology and Oncology, Leesburg, FL, USA; ¹⁰Minnesota Oncology Clinic, Medical Oncology and Hematology, Burnsville, MN, USA; ¹¹BeiGene (Beijing) Co, Ltd, Beijing, China, and BeiGene USA, Inc, Hematology Clinical Development, San Mateo, CA, USA; ¹²Willamette Valley Cancer Institute and Research Center, Medical Oncology and Hematology, Eugene, OR, USA

INTRODUCTION

- Bruton tyrosine kinase (BTK) inhibitors are a mainstay of treatment for B-cell malignancies; however, their use can be limited by adverse events (AEs), many of which are potentially caused by off-target inhibition of other tyrosine kinases^{1,3}
- Zanubrutinib is a potent and selective next-generation BTK inhibitor designed to maximize BTK occupancy and minimize off-target kinase binding and associated AEs⁴
- Previous results from an ongoing phase 2 study (BGB-3111-215; NCT04116437) showed that zanubrutinib is well tolerated in patients who are intolerant of ibrutinib and/or acalabrutinib⁵
- Zanubrutinib demonstrated higher selectivity than ibrutinib, acalabrutinib, and acalabrutinib's major metabolite, M27, by kinase profiling (Figure 1)^{5,6}
- Zanubrutinib, ibrutinib, acalabrutinib, and M27 (metabolite of acalabrutinib) demonstrated >50% inhibition of 7, 17, 15, and 23 kinases, respectively, of the 370 kinases tested
- Here, we report updated results on the tolerability and efficacy of zanubrutinib in patients intolerant of acalabrutinib (Cohort 2)

Figure 1. Kinase Selectivity of Zanubrutinib, Ibrutinib, Acalabrutinib, and Acalabrutinib Metabolite M27




Reprinted from Shadman M, et al. *Lancet Haematol*. 2023;10(10):e35-e45. Copyright © 2022 Elsevier Ltd.

METHODS

- BGB-3111-215 is an ongoing phase 2 study (Figure 2) in patients with previously treated B-cell malignancies who were intolerant of acalabrutinib, as defined by 1 of the following:
 - Grade ≥1 nonhematologic toxicities with ≥3 recurrent episodes or lasting >7 days, or grade ≥3 of any duration
 - Grade ≥3 febrile neutropenia of any duration
 - Grade 4 heme toxicity that persisted to the point that the investigator chose to stop therapy due to toxicity, not progression
 - Inability to use acid-reducing agents or anticoagulants (eg, proton pump inhibitors, warfarin) due to concurrent acalabrutinib use
- Patients with Richter transformation or progressive disease (PD) while on prior BTK inhibitor treatment were excluded

Figure 2. BGB-3111-215 Study Design



ClinicalTrials.gov: NCT04116437; data cutoff: May 15, 2023
 * Study is ongoing.
 CLL, chronic lymphocytic leukemia; DCR, disease control rate; MCL, mantle cell lymphoma; MZL, marginal zone lymphoma; SLL, small lymphocytic lymphoma; WM, Waldenström macroglobulinemia.

RESULTS

Patients

- As of May 15, 2023, 27 acalabrutinib-intolerant patients had enrolled (Table 1); 13 of these patients were also intolerant of ibrutinib

Table 1. Patient Demographics and Baseline Characteristics

Characteristic	Acalabrutinib-Intolerant (n=27)
Indication, n (%)	
CLL	17 (63)
SLL	2 (7)
MCL	2 (7)
MZL	2 (7)
WM	4 (15)
Age, median (range), years	73 (51-87)
Sex, n (%)	
Male	17 (63)
Female	10 (37)
ECOG PS, n (%)	
0	18 (67)
1	7 (26)
2	2 (7)
No. of prior anticancer therapy regimens, median (range)	2 (1-6)
Prior BTK inhibitor, n (%)	
Ibrutinib monotherapy	12 (44)
Ibrutinib combination therapy	1 (4)
Acalabrutinib monotherapy	26 (96)
Acalabrutinib combination therapy	1 (4)
Cumulative acalabrutinib exposure, median (range), months	5.4 (0.5-33.7)
On-study zanubrutinib dosing regimen, n (%)	
160 mg BID	19 (70)
320 mg QD	8 (30)

CLL, chronic lymphocytic leukemia; ECOG PS, Eastern Cooperative Oncology Group performance status; MCL, mantle cell lymphoma; MZL, marginal zone lymphoma; SLL, small lymphocytic lymphoma; WM, Waldenström macroglobulinemia.

- Of 27 acalabrutinib-intolerant patients, 7 (26%) discontinued zanubrutinib treatment (Table 2) (AE, n=2; physician decision, n=2; withdrawal by patient, n=2; PD, n=1)

Table 2. Patient Disposition


Patients, n (%)	Acalabrutinib-Intolerant (n=27)
Remaining on treatment	20 (74)
Remaining on study	23 (85)
Discontinued from treatment	7 (26)
AE	2 (7) ^a
Physician decision	2 (7)
Withdrawal by patient	2 (7)
PD	1 (4)
Death, n (%)	1 (4)
Zanubrutinib treatment duration, median (range), months	11.4 (0.5-32.2)
Survival follow-up, median (range), months	12.4 (1.6-32.2)

^a Myalgia (n=1), diarrhea (n=1).

Safety

- Of 40 acalabrutinib-intolerance events that were reported by 27 patients, the most common (≥2 events) were arthralgia (n=6), headache (n=5), myalgia (n=5), diarrhea (n=3), rash (n=3), fatigue (n=2), and hemorrhage (n=2) (Figure 3)
- Most acalabrutinib-intolerance events (28 of 40; 70%) did not recur at any grade with zanubrutinib; of the 12 that did recur, none recurred at a higher severity
- Seventeen of 27 patients (63%) did not experience any recurrence of their prior acalabrutinib-intolerance events
- Two patients discontinued zanubrutinib due to recurrence of a prior acalabrutinib-intolerance event (grade 2 myalgia, n=1; and grade 3 diarrhea, n=1; both recurred at the same grade)
- Three of 27 patients (11%) experienced the same intolerance event (pain in extremity, diarrhea, and atrial fibrillation; n=1 each) with ibrutinib and acalabrutinib
 - Two (67%) did not have a recurrence of those events with zanubrutinib
 - One (33%) had a recurrence at a lower grade (diarrhea)

Figure 3. Recurrence of Acalabrutinib-Intolerance Events on Zanubrutinib



* No events recurred at a higher grade.

- No AEs led to death, and 2 events (7%) led to treatment discontinuation (Table 3)
- The most common grade ≥3 AE was neutrophil count decreased, which occurred in 3 patients (11%) (Table 4)
- Anemia and thrombocytopenia/platelet count decreased did not occur

Table 3. Adverse Event Summary

Patients, n (%)	Any Grade (n=27)
Serious AE	7 (26)
Leading to treatment discontinuation	2 (7)
Leading to dose interruption	16 (59)
Leading to dose reduction	6 (22)
Leading to death	0

CONCLUSIONS

- The median zanubrutinib exposure was 6 months longer than the reported cumulative acalabrutinib exposure before discontinuation (11.4 months vs 5.4 months, respectively)
- Most patients (63%) did not experience any recurrence of their prior acalabrutinib-intolerance event
- Of the 40 acalabrutinib-intolerance events, 28 did not recur; of the 12 that did recur, none recurred at a higher grade
- Zanubrutinib provided clinically meaningful benefit in efficacy-evaluable patients who were previously intolerant of acalabrutinib, as measured by a disease control rate of 96%
- The results from this study demonstrate that zanubrutinib may be a viable treatment option for patients who are intolerant of acalabrutinib

Table 4. Select Adverse Events^a


Patients, n (%)	Any Grade (n=27)	Grade ≥3 (n=27)
Any AE	26 (96)	12 (44) ^b
Neutrophil count decreased	3 (11)	3 (11)
Neutropenia	2 (7)	2 (7)
Diarrhea	12 (44)	1 (4)
Hypertension	6 (22)	1 (4)
COVID-19	5 (19)	1 (4)
Maculopapular rash	3 (11)	1 (4)
Abdominal pain	2 (7)	1 (4)
Bacteremia	1 (4)	1 (4)
Cellulitis	1 (4)	1 (4)
COVID-19 pneumonia	1 (4)	1 (4)
Fall	1 (4)	1 (4)
Febrile neutropenia	1 (4)	1 (4)
Gastroenteritis salmonella	1 (4)	1 (4)
Hip fracture	1 (4)	1 (4)
Pneumonia	1 (4)	1 (4)
Small intestinal obstruction	1 (4)	1 (4)

^a AEs shown in this table occurred in at least 1 patient at grade ≥3 severity; any-grade data for these select AEs are also shown. ^b Some patients had more than 1 grade ≥3 event.

Efficacy

- Among the 25 efficacy-evaluable patients on zanubrutinib, 24 (96%) achieved SD or better, and 16 (64%) achieved minor response (MR) or better (Figure 4)
- Twelve of 17 efficacy-evaluable patients (71%) with CLL/SLL on zanubrutinib achieved a partial response with lymphocytosis (PR-L) or better

Figure 4. BOR by Investigator Assessment



^a In patients with a BOR better than SD. BOR, best overall response; DCR, disease control rate; MR, minor response; PR-L, partial response with lymphocytosis.

REFERENCES

- Stephens DM, Byrd JC. *Blood*. 2019;133(12):1298-1307.
- Furman RR, et al. *Leukemia*. 2021;35(11):3201-3211.
- Mato AR, et al. *Haematol*. 2018;103(5):874-879.
- Guo Y, et al. *J Med Chem*. 2019;62(17):7923-7940.
- Shadman M, et al. *Lancet Haematol*. 2023;10(11):e35-e45.
- Shadman M, et al. *Blood*. 2021;138(suppl 1):1410.

ACKNOWLEDGMENTS

The authors would like to thank the investigators, site support staff, and especially the patients for participating in this study. This study was sponsored by BeiGene, Ltd. Medical writing and editorial assistance was provided by Nucleus Global, an Inizio company, and supported by BeiGene.