

First-Line Tislelizumab Plus Chemotherapy in Gastric/Gastroesophageal Junction Cancer: RATIONALE-305 Asian Subgroup

Authors: Ken Kato,¹ Yuxian Bai,² Jianhua Shi,³ Keun-wook Lee,⁴ Jufeng Wang,⁵ Hongming Pan,⁶ Sun Young Rha,⁷ Ruixing Zhang,⁸ Hidekazu Hirano,¹ Kensei Yamaguchi,⁹ Zengqing Guo,¹⁰ Yi Ba,¹¹ Lei Yang,¹² Hiroshi Tsukuda,¹³ Yaling Xu,¹⁴ Tao Sheng,¹⁵ Silu Yang,¹⁴ Liyun Li,¹⁴ Do-Youn Oh,¹⁶ Rui-hua Xu,¹⁷

Affiliations: ¹National Cancer Center Hospital, Tokyo, Japan; ²Harbin Medical University Cancer Hospital, Harbin, China; ³Linyi Cancer Hospital, Linyi, China; ⁴Seoul National University College of Medicine, Seoul National University Bundang Hospital, Seongnam, Republic of Korea; ⁵Henan Cancer Hospital, Zhengzhou, China; ⁶Sir Run Run Shaw Hospital, Zhejiang University School of Medicine, Hangzhou, China; ⁷Yonsei Cancer Center, Yonsei University Health System, Seoul, Republic of Korea; ⁸Fourth Hospital of Hebei Medical University, Shijiazhuang, China; ⁹Cancer Institute Hospital of JFCR, Tokyo, Japan; ¹⁰Fujian Cancer Hospital, Fuzhou, China; ¹¹Tianjin Medical University Cancer Institute & Hospital, Tianjin, China; ¹²Nantong Cancer Hospital, Nantong, China; ¹³Izumi Municipal Hospital, Izumi, Japan; ¹⁴BeiGene (Beijing) Co., Ltd, Beijing, China; ¹⁵BeiGene, Ltd, Boston, MA, United States; ¹⁶Seoul National University Hospital, Cancer Research Institute, Seoul National University College of Medicine, Seoul, Republic of Korea; ¹⁷Sun Yat-sen University Cancer Center State Key Laboratory of Oncology in South China, Collaborative Innovation Center of Cancer Medicine, Guangzhou, China.

ABSTRACT

Background: Survival outcomes with platinum + fluoropyrimidine chemotherapy (chemo) alone as first-line (1L) therapy for advanced gastric/gastroesophageal junction cancer (GC/GEJC) are poor. Tislelizumab (TIS), an anti-PD-1 antibody, + chemo, showed significant overall survival (OS) benefit vs placebo (PBO) + chemo as 1L therapy in patients (pts) with advanced GC/GEJC in the randomized, double-blind, global phase 3 RATIONALE-305 study (NCT03777657). Here, we present results from the Asian pt subgroup analysis.

Methods: Eligible pts were randomized (1:1) to receive TIS 200 mg intravenously every 3 weeks plus investigator-chosen chemo (ICC [XELOX or FP]) or PBO + ICC until disease progression, or intolerable toxicity. Dual primary endpoints were OS in both the programmed death ligand-1 (PD-L1)+ (tumor area positivity $\geq 5\%$) and ITT analysis sets. Secondary endpoints included progression-free survival, objective response rate, duration of response, and safety.

Results: 748 pts (TIS + chemo: n=376, PBO + chemo: n=372, of N=997) enrolled from Asia (China, Korea and Japan) were analyzed, including 403 pts in the PD-L1+ analysis set. At final analysis, TIS + chemo showed improvements vs PBO + chemo in OS both in PD-L1+ (median [m] OS: 18.0 [95% confidence interval (CI): 15.0, 21.6] vs 14.0 mo [95% CI: 12.3, 15.5], hazard ratio [HR]=0.71 [95% CI: 0.56, 0.89]) and ITT analysis set (mOS: 16.4 [95% CI: 14.4, 18.0] vs 14.1 mo [95% CI: 12.8, 15.4], HR=0.83 [95% CI: 0.70, 0.97]). Incidences of grade ≥ 3 treatment-related adverse events (TRAEs) were 55.5% vs 50.0% and TRAEs leading to death were 2.4% vs 0.8% for TIS vs PBO, respectively. TRAEs leading to discontinuation were reported in 17.1% pts in TIS + chemo arm vs 8.9% patients in PBO + chemo arm.

Conclusions: TIS + chemo produced a clinically meaningful improvement in OS vs PBO + chemo and a safety profile in pts with advanced GC/GEJC in the Asian subgroup that was consistent with published results in the overall study population.